



## Last glacial–interglacial vegetation and environmental dynamics in southern Siberia: Chronology, forcing and feedbacks

Elena V. Bezrukova<sup>a</sup>, Pavel E. Tarasov<sup>b,\*</sup>, Nadia Solovieva<sup>c</sup>, Sergey K. Krivonogov<sup>d</sup>, Frank Riedel<sup>b</sup>

<sup>a</sup> Institute of Geochemistry, Siberian Branch Russian Academy of Sciences, Favorsky Str. 1a, Irkutsk, 664033, Russia

<sup>b</sup> Institute of Geological Sciences, Palaeontology, Free University, Malteserstr. 74–100, House D, Berlin, 12249, Germany

<sup>c</sup> Department of Geography, University College London, Gower Str., London WC1E 6BT, UK

<sup>d</sup> Institute of Geology and Mineralogy, Russian Academy of Sciences, Siberian Branch, Koptug ave. 3, Novosibirsk, 630090, Russia

### ARTICLE INFO

#### Article history:

Received 15 January 2010

Received in revised form 18 July 2010

Accepted 21 July 2010

Available online 25 July 2010

#### Keywords:

Eurasia

Pollen and diatom records

Quantitative vegetation reconstruction

Late Pleistocene interstadials

Last glacial maximum

Younger Dryas

Holocene climatic optimum

### ABSTRACT

Radiocarbon-dated pollen and diatom records from Lake Kotokel in southern Siberia help to reconstruct the environmental history of the area since ~47 kyr BP. Pollen spectra composition and reconstructed biome scores suggest predominance of a tundra–steppe vegetation and variable woody cover (5–20%) between ~47 and 30 kyr BP, indicating generally a harsh and unstable climate during this interval, conventionally regarded as the interstadial within the last glacial. The short-term climate amelioration episodes in the glacial part of the records are marked by the peaks in taiga and corresponding minima in steppe biome scores and appear synchronously with the hemispheric temperature and precipitation changes recorded in the Greenland ice cores and Chinese stalagmites. Transition to full glacial environments occurred between 32 and 30 kyr BP. The interval at ~30–24 kyr BP was probably the driest and coldest of the whole record, as indicated by highest scores for steppe biome, woody coverage <5%, absence of diatoms and reduced size of the lake. A slight amelioration of the regional climate at ~24–22 kyr BP was followed by a shorter than the previous and less pronounced deterioration phase. The late-glacial (~17–11.65 kyr BP) is marked by a gradual increase in tree/shrub pollen percentages and re-appearance of diatoms. After 14.7 kyr BP the climate became warmer and wetter than ever during ~47–14.7 kyr BP, resulting in the deepening of the lake and increase in the woody coverage to 20–30% ~14.5–14 kyr and ~13.3–12.8 kyr BP. These two intervals correspond to the Meindorf and Allerød interstadials, which until now were interpreted as part of the undifferentiated Bölling/Allerød interstadial complex in the Lake Baikal region. The increase in tundra biome scores and pronounced change in the diatom composition allow (for the first time) the unambiguous identification of the Younger Dryas (YD) in the Lake Baikal region at ~12.7–11.65 kyr BP, in agreement with the formal definition and dating of the YD based on the Greenland NGRIP ice core records. The maximal spread of the taiga communities in the region is associated with a warmer and wetter climate than the present prior to ~7 kyr BP. This was followed by a wide spread of Scots pine, indicating the onset of modern environments.

© 2010 Elsevier B.V. All rights reserved.

### 1. Introduction

Worldwide terrestrial and marine sedimentary archives demonstrate that the last 50-kyr interval in the Earth's history experienced a number of long- and short-term climatic oscillations (Guiot et al., 1989; Vincens et al., 2005; Heusser et al., 2006; Bout-Roumazilles et al., 2007; Svensson et al., 2008). High-resolution and accurately dated pollen and sedimentary records of the late-glacial/early Holocene interval exist for several regions of Europe (Litt and Stebich, 1999; Allen and Huntley, 2000; Brauer et al., 2008) and East Asia (Stebich

et al., 2009), providing important insight into the environmental dynamics in the North Pacific and North Atlantic regions. However, a recent global-scale synthesis of the Holocene climatic data (Wanner et al., 2008) demonstrates a lack of palaeorecords of comparable dating quality and resolution from the vast areas of Eurasia, including Siberia and Central Asia. The dating problem becomes even more obvious, when the pre-Holocene interval of the late Quaternary is considered (Rudaya et al., 2009; Shichi et al., 2009).

Southern Siberia – the region of Russia between ~80–120°E and ~50–60°N – consists of numerous sub-latitudinal mountain ranges and lakes, including Lake Baikal in the east (Fig. 1a and b). The lake sediments are one of the most promising sources of detailed palaeoenvironmental information, which provide an opportunity for bridging the European and Asian palaeoclimate archives and addressing critical questions concerning Quaternary climatology, e.g. reconstructing atmospheric circulation patterns (Williams et al.,

\* Corresponding author. Tel.: +49 30 83870280; fax: +49 30 83870745.

E-mail addresses: [bezrukova@igc.irk.ru](mailto:bezrukova@igc.irk.ru) (E.V. Bezrukova),

[ptarasov@zedat.fu-berlin.de](mailto:ptarasov@zedat.fu-berlin.de) (P.E. Tarasov), [nsoloviev@geog.ucl.ac.uk](mailto:nsoloviev@geog.ucl.ac.uk) (N. Solovieva),

[paleobio@zedat.fu-berlin.de](mailto:paleobio@zedat.fu-berlin.de) (F. Riedel).